

THE ZERO-TYPE PROPERTY AND MIXING OF BERNOULLI SHIFTS

ZEMER KOSLOFF

ABSTRACT. We prove that every non-singular Bernoulli shift is either zero-type or there is an equivalent invariant stationary product probability. We also give examples of a type III_1 Bernoulli shift and a Markovian flow which are power weakly mixing and zero type.

Introduction. This article deals with the concept of zero-type for invertible non-singular transformations T of the standard probability space (X, \mathcal{B}, m) .

A non-singular transformation without an absolutely continuous invariant probability is called zero-type, sometimes also called mixing, if its Koopman operator is mixing, meaning that it's maximal spectral type is a Rajchman measure. This generalises the notion of zero-type introduced by Hajian and Kakutani [HK]. The name mixing comes from the fact that in the case of probability preserving transformations the classical notion of mixing is equivalent to mixing of the Koopman operator restricted to $L_0^2(X, m) = L^2(X, m) \ominus \mathbb{C}$. In the interplay between non singular ergodic theory and infinitely divisible processes, the zero type transformations are associated with the classical notion of mixing of infinitely divisible processes, see [RS] or [Ro].

In the first part we prove that every non-singular Bernoulli shift is either zero-type or there is an equivalent invariant product probability. Thus the Hamachi shift [Ham] and the type III_1 shift given in [Kos] are examples of conservative, ergodic, zero-type transformations which do not posses an m equivalent σ -finite invariant measure, also known as type III and zero type transformations. For another construction of zero type, type III transformations see [Da1, Theorem 0.3].

A non singular transformation T is weakly mixing if $S \times T$ is ergodic for every ergodic probability preserving transformation S . It follows that if T is a probability preserving weakly mixing transformation then $T \times \cdots \times T$ (k -times) is ergodic for every $k \in \mathbb{N}$. In general this is not true, as there exist weakly mixing transformations such that $T \times T$ is not ergodic. In an attempt to understand the notion of weak mixing for non singular transformations the concept of power weakly mixing transformations was introduced. T is called *power weakly mixing* if for any $l_1, l_2, \dots, l_k \in \mathbb{Z} \setminus \{0\}$,

$$(X^k, P^{\otimes k}, T^{l_1} \times T^{l_2} \times \cdots \times T^{l_k})$$

This research was supported by THE ISRAEL SCIENCE FOUNDATION grant No. 1114/08.

is an ergodic automorphism where

$$P^{\otimes k} = \underbrace{P \otimes P \otimes \cdots \otimes P}_{k\text{-times}}.$$

A weaker notion is that of infinite ergodic index which means that for every $k \in \mathbb{N}$, the k -fold product of T is an ergodic automorphism. In [AFS] it was shown that Chacon's non singular type III_λ transformation, $0 < \lambda \leq 1$, is a power weakly mixing transformation. In [Da2] a construction of a transformation which is of infinite ergodic index but not power weakly mixing was given. Later in [DP] a flow was constructed such that all times are of infinite ergodic index. These constructions use the cutting and stacking method which usually doesn't give a zero type transformation.

We give a construction of a Bernoulli shift which is power weakly mixing and type III_1 . By the first part it is also zero type. Finally a continuous time flow $\{\phi_t\}_{t \in \mathbb{R}}$ is given such that for every $a_1, a_2, \dots, a_n \in \mathbb{R}$,

$$\phi_{a_1} \times \phi_{a_2} \times \cdots \times \phi_{a_n}$$

is ergodic. This flow is the time shifts of a continuous time Markov Chain.

Preliminaries: Let (X, \mathcal{B}, P) be a probability space. An invertible measurable transformation $T : X \rightarrow X$ is said to be *non-singular* with respect to P if it preserves the P -null sets. i.e for every $A \in \mathcal{B}$, $P(A) = 0$ if and only if

$$P \circ T(A) := P(TA) = 0.$$

A measure m on X is said to be *T -invariant* if for every $A \in \mathcal{B}$,

$$P \circ T(A) = P(A).$$

If T is non-singular, then for every n , $P \circ T^n$ is absolutely continuous with respect to P . By the Radon-Nikodym theorem there exist measurable functions $\frac{dP \circ T^n}{dP} \in L_1(X, P)_+$ such that for every $A \in \mathcal{B}$,

$$P \circ T^n(A) = \int_A \left(\frac{dP \circ T^n}{dP} \right) dP.$$

Denote $(T^n)'(x) := \frac{dP \circ T^n}{dP}$.

A set $W \in \mathcal{B}$ is called *wandering* if $\{T^n W\}_{n=-\infty}^{\infty}$ are disjoint. As in [Aar, p.7] denote by \mathfrak{D} the measurable union of all wandering sets for T , this set is T -invariant. Its complement is denoted by \mathfrak{C} .

Definition. We call (X, \mathcal{B}, P, T) *dissipative* if $\mathfrak{D} = X$. If $\mathfrak{C} = X$ then (X, \mathcal{B}, P, T) is said to be *conservative*.

We will use the following version of Hopf's decomposition theorem which says that the conservative and dissipative parts can be separated in the following way.

Theorem I. *For every non-singular transformation T of the probability space (X, \mathcal{B}, P) there exists a decomposition $X = \mathfrak{D} \cup \mathfrak{C}$ such that $(X, \mathcal{B}, P|_{\mathfrak{D}}, T)$ is dissipative and $(X, \mathcal{B}, P|_{\mathfrak{C}}, T)$ is conservative. Furthermore*

$$\sum_{k=0}^{\infty} (T^{-k})'(x) < \infty \text{ a.e. } x \in \mathfrak{D}$$

and

$$\sum_{k=0}^{\infty} (T^{-k})'(x) = \infty \text{ a.e. } x \in \mathfrak{C}.$$

The Hellinger Integral and definition of the Zero Type Property. The Koopman operator $U : L_2(X, \mathcal{B}, P)$ is then defined by

$$Uf(x) = \sqrt{T'(x)} f \circ T(x).$$

It is a unitary operator and by the chain rule for the Radon-Nikodym derivatives for every $n \in \mathbb{Z}$,

$$U^n f = \sqrt{(T^n)'} f \circ T^n.$$

Definition. A transformation is *Non-Singular zero-type* (NS zero type) if the maximal spectral type $\sigma_T \in \mathcal{P}(\mathbb{T}, \mathcal{B}(\mathbb{T}))$ of U_T is a Rajchman measure. That is its Fourier coefficients $\hat{\sigma}_T(n)$ tend to 0 as $n \rightarrow \infty$. and so for every $f \in L_2(X, P)$,

$$\int_X f \cdot U^n f dP \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Remark. By looking at the Koopman operators it is seen that the zero-type property depends only on the equivalence class of P .

Remark. In the case when (X, \mathcal{B}, m, T) is a σ -finite measure preserving transformation Krenzel and Sucheston [KrS] have showed that mixing of the Koopman operator is equivalent to the Hajian and Kakutani zero-type condition [HK] which states that for every $A \in \mathcal{B}$ with $m(A) < \infty$,

$$\lim_{n \rightarrow \infty} m(A \cap T^{-n}A) = 0.$$

Definition. Let P, Q be two probability measures on X . The *Hellinger Integral* (see [Hel] or [Kak]) is then defined as

$$\rho(P, Q) = \int_X \sqrt{\frac{dP}{dm}} \cdot \sqrt{\frac{dQ}{dm}} dm$$

where m is any finite measure on X such that $P \ll m$, $Q \ll m$. In the special case where $P \ll Q$ we can take $m = Q$ and have

$$\rho(P, Q) = \int_X \sqrt{\frac{dP}{dQ}} dQ.$$

The function $\rho(\cdot, \cdot)$ measures the amount of singularity of P with respect to Q . This function satisfies that for every $P, Q \in \mathcal{P}(X)$, $0 \leq \rho(P, Q) \leq 1$. Also $\rho(P, Q) = 0$ if and only if P is singular with respect to Q .

The proof of the following proposition is standard.

Proposition 1. *Let T be a non-singular transformation of the probability space (X, \mathcal{B}, P) . The following are equivalent.*

- (i) $\lim_{n \rightarrow \infty} \rho(P, P \circ T^n) = 0$.
- (ii) $(T^n)' \xrightarrow{P} 0$.
- (iii) σ_T is a Rajchman measure.

1. BERNOULLI SHIFTS ARE ZERO-TYPE OR MIXING

Let $\mathbb{X} = \{0, 1\}^{\mathbb{Z}}$ and T be the left shift action on \mathbb{X} , that is

$$(Tw)_i = w_{i+1}.$$

Denote the cylinder sets by

$$[b]_k^l = \{w \in \mathbb{X} : \forall i = k, \dots, l, w_i = b_i\}.$$

A measure $P = \prod_{k=-\infty}^{\infty} P_k \in \mathcal{P}(\mathbb{X})$ is called a product measure if for every $k < l$, and for every cylinder $[b]_k^l$,

$$P([b]_k^l) = \prod_{j=k}^l P_j(\{b_j\}).$$

We will say that a product measure P is *non-singular* if the shift is non-singular with respect to P .

For two product probability measures P, Q and $N \in \mathbb{N} \cup \{\infty\}$ define

$$d_N(P, Q) := \sum_{k=-N}^N \left\{ \left(\sqrt{P_k(\{0\})} - \sqrt{Q_k(\{0\})} \right)^2 + \left(\sqrt{P_k(\{1\})} - \sqrt{Q_k(\{1\})} \right)^2 \right\}.$$

Notice that $d_N(P, Q) \uparrow d_\infty(P, Q)$ as $N \rightarrow \infty$. Set $d(P, Q) := d_\infty(P, Q)$

The following lemma is a direct consequence of Kakutani's Theorem on equivalence of product measures[Kak].

Lemma 2. *Let $P = \prod_{k=-\infty}^{\infty} P_k$ be a product measure. Then*

(1) *For any two product measures P, Q ,*

$$d(P, Q) \propto -\log \rho(P, Q).$$

(2) *P is non-singular if and only if*

$$(1.1) \quad d(P, P \circ T) < \infty.$$

(3) *The shift is NS zero-type if and only if*

$$\lim_{n \rightarrow \infty} d(P, P \circ T^n) = \infty.$$

Theorem 3. *Let $P = \prod_{k=-\infty}^{\infty} P_k$ be a non-singular product measure. Then either there exists a shift invariant P -equivalent probability or the shift $(\mathbb{X}, \mathcal{B}(\mathbb{X}), P, T)$ is NS zero-type. Therefore a non-singular shift is either mixing in the probability preserving sense or mixing in the non-singular sense.*

Theorem 1 follows from lemmas 4 and 5.

Lemma 4. *Let P be a non-singular product measure on $\{0, 1\}^{\mathbb{Z}}$ such that*

$$\exists \lim_{k \rightarrow -\infty} P_k = (p, 1-p) := \mu_p.$$

Denote by $Q = \prod_{k=-\infty}^{\infty} \mu_p$. Then if $P \perp Q$ then $(\mathbb{X}, \mathcal{B}(\mathbb{X}), P, T)$ is of NS-zero type. Else Q is a P -equivalent shift-invariant probability measure.

Proof. Assume that $P \perp Q$. Then by Kakutani's theorem

$$d(P, Q) = \infty.$$

By claim 2 its enough to show that $\lim_{n \rightarrow \infty} d(P, P \circ T^n) = \infty$.

Let $M > 0$. Since $d(P, Q) = \infty$ there exists a $N \in \mathbb{N}$ such that

$$d_N(P, Q) > M.$$

For every $n \in \mathbb{N}$,

$$\begin{aligned} d(P, P \circ T^n) &\geq \sum_{k=-N}^N \left\{ \left(\sqrt{P_k(\{0\})} - \sqrt{P_{k-n}(\{0\})} \right)^2 \right. \\ &\quad \left. + \left(\sqrt{P_k(\{1\})} - \sqrt{P_{k-n}(\{1\})} \right)^2 \right\} \end{aligned}$$

Therefore since $\lim_{j \rightarrow -\infty} P_j = \mu_p$ then

$$\begin{aligned} \liminf_{n \rightarrow \infty} d(P, P \circ T^n) &\geq d_N(P, Q) \\ &\geq M. \end{aligned}$$

Since M is arbitrary then

$$\lim_{n \rightarrow \infty} d(P, P \circ T^n) = \infty.$$

□

Lemma 5. *Let P be a non-singular product measure on $\{0, 1\}^{\mathbb{Z}}$ such that*

$$\liminf_{k \rightarrow -\infty} P_k(\{0\}) \neq \limsup_{k \rightarrow -\infty} P_k(\{0\}).$$

Then $(\mathbb{X}, \mathcal{B}(\mathbb{X}), P, T)$ is NS zero-type.

Proof. Write $q_1 = \lim_{k \rightarrow -\infty} \inf P_k(\{0\})$ and $q_2 = \lim_{k \rightarrow -\infty} \sup P_k(\{0\})$.

Let $M > 0$. Set $\alpha = \frac{q_2 - q_1}{4}$. Define

$$A_{q_i} := \{n \in \mathbb{Z} : |P_n(\{0\}) - q_i| < \alpha\}, \quad i = 1, 2.$$

Let $A_{q_i}^N = A_{q_i} \cap [-N, N]$.

Choose N large enough so that

$$|A_q^N| \geq \frac{M}{\alpha} \quad \text{and} \quad |A_p^N| \geq \frac{M}{\alpha}.$$

Since $d(P, P \circ T) < \infty$ then for every $j \in \mathbb{Z} \cap [-N, N]$,

$$\lim_{n \rightarrow \infty} |P_{-n}(\{0\}) - P_{-n+j}(\{0\})| = 0.$$

Therefore for large enough $n \in \mathbb{N}$ either

$$[-N - n, N - n] \cap A_{q_1} = \emptyset$$

or

$$[-N - n, N - n] \cap A_{q_2} = \emptyset.$$

Therefore for large enough $n \in \mathbb{N}$,

$$\begin{aligned} d(P, P \circ T^n) &\geq d_N(P, Q) \\ &\geq \sum_{k \in A_{q_1}^N} \left(\sqrt{P_k(\{0\})} - \sqrt{P_{k-n}(\{0\})} \right)^2 \\ &\quad + \sum_{k \in A_{q_2}^N} \left(\sqrt{P_k(\{0\})} - \sqrt{P_{k-n}(\{0\})} \right)^2 \\ &\geq \min(\alpha \cdot |A_{q_1}^N|, \alpha \cdot |A_{q_2}^N|) \geq M. \end{aligned}$$

Therefore

$$\lim_{n \rightarrow \infty} d(P, P \circ T^n) = \infty$$

and the shift is NS zero-type. □

2. A ZERO TYPE AND POWER WEAKLY MIXING BERNOULLI SHIFT

In this section we construct a Bernoulli shift which is zero type and power weak mixing. The construction is done by imposing a stronger growth condition on the shift constructed in [Kos].

Construction of the product measure:

The product measure will be $P = \prod_{k=-\infty}^{\infty} P_k$, where

$$(2.1) \quad \forall i \geq 0, P_i(0) = P_i(1) = \frac{1}{2}.$$

The definition of P_k for negative k 's is more complicated as it involves an inductive procedure.

2.1. The inductive definition of P_k for negative k 's. We will need to define inductively 5 sequences $\{\lambda_t\}_{t=1}^{\infty}$, $\{n_t\}_{t=1}^{\infty}$, $\{m_t\}_{t=1}^{\infty}$, $\{M_t\}_{t=0}^{\infty}$ and $\{N_t\}_{t=1}^{\infty}$. The sequence $\{\lambda_t\}$ is of real numbers which decreases to 1. The other four, $\{n_t\}_{t=1}^{\infty}$, $\{m_t\}_{t=1}^{\infty}$, $\{M_t\}_{t=0}^{\infty}$ and $\{N_t\}_{t=1}^{\infty}$ are increasing sequences of integers.

First choose a positive summable sequence $\{\epsilon_t\}_{t=1}^{\infty}$ and set $M_0 = 1$.

Base of the induction: Set $\lambda_1 = 2$, $n_1 = 2$, $m_1 = 4$. Set also $N_1 = M_0 + n_1 = 3$ and $M_1 = N_1 + m_1 = 7$.

Given $\{\lambda_u, n_u, N_u, m_u, M_u\}_{u=1}^{t-1}$, we will choose the next level $\{\lambda_t, n_t, N_t, m_t, M_t\}$ in the following order. First we choose λ_t depending on M_{t-1} and ϵ_t . Given λ_t we will choose n_t and then N_t will be defined by

$$N_t := M_{t-1} + n_t.$$

Then given N_t we will choose m_t and finally set

$$M_t := N_t + m_t.$$

Choice of λ_t : Set $k_t := \left\lfloor \log_2 \left(\frac{M_{t-1}}{\epsilon_t} \right) \right\rfloor + 1$, where $\lfloor x \rfloor$ denotes the integral part of x . Then set $\lambda_t = e^{\frac{1}{2^{k_t}}}$. With this choice of λ_t we have,

$$(2.2) \quad \lambda_t^{M_{t-1}} < e^{\epsilon_t}.$$

This choice of λ_t has the property that for every $u < t$, $\lambda_u = \lambda_t^{2^{k_t - k_u}}$.

Define

$$A_{t-1} := \left\{ \prod_{u=1}^{t-1} \lambda_u^{x_u} : x_u \in [-n_u, n_u] \right\}.$$

Choice of n_t : Given $\{\lambda_u, n_u, m_u\}_{u=1}^{t-1}$ and λ_t , the set A_{t-1} is a finite subset of $\lambda_t^{\mathbb{Z}}$. Choose n_t which satisfies

$$\lambda_t^{n_t/4} \geq \max \{a^2 : a \in A_{t-1}\}.$$

Choice of m_t : Now that n_t is chosen we set $N_t = M_{t-1} + n_t$. Set

$$(2.3) \quad tN_t (2 + 2^{tN_t}) = m_t.$$

Remark. Since m_t satisfies 2.3 then for every $k < t$ and $n \leq N_t$,

$$(2.4) \quad \frac{m_t}{n} - N_t > 2^{kN_t}$$

Definition 6. Let (X, μ, T) be a non singular automorphism such that $\mathcal{B}(X) \neq \{\emptyset, X\}$ and let $\mathcal{F} \subset \mathcal{B}$ be a factor algebra. Then:

- (i) \mathcal{F} is *exhaustive* if $\bigvee_{n=0}^{\infty} T^n \mathcal{F} = \mathcal{B}$.
- (ii) \mathcal{F} is *exact* if $\bigcap_{n=0}^{\infty} T^n \mathcal{F} = \{\emptyset, X\}$.
- (iii) T is a *K-automorphism* if it is conservative and admits a factor algebra $\mathcal{F} \subset \mathcal{B}$ that is exhaustive and exact and such that T' is \mathcal{F} measurable.

Remark. Krengel has shown in [Kre, p. 153-154] that all K -automorphisms are ergodic. See also [ST, Proposition 4.8(a)].

Theorem 7. *The Bernoulli shift $(\mathbb{X}, \mathcal{B}(\mathbb{X}), P, T)$ is a non-singular, type III₁, zero-type and power weakly mixing transformation.*

Proof. In [Kos] it is shown that the shift is a type III₁ transformation. Therefore by Theorem 3 the shift is a zero type transformation. It remains to show that the shift is power weak mixing.

Let $l_1, l_2, \dots, l_k \in \mathbb{Z} \setminus \{0\}$ and denote by $S := T^{l_1} \times T^{l_2} \times \dots \times T^{l_k}$. Clearly

$$S^{n'}(w_1, w_2, \dots, w_k) = \prod_{i=1}^k T^{(l_i n)'}(w_i)$$

and $(\mathbb{X}^k, P^{\times k}, S)$ admits an exhaustive and exact factor. Therefore in order to prove the ergodicity of S it is sufficient to show that S is conservative.

By [Kos, lemma 3] and a similar calculation for negative n 's there exists $t_0 \in \mathbb{N}$ such that for every $t > t_0$, $|n| \in [N_t, m_t)$ and $w \in \mathbb{X}$

$$T^{n'}(w) \geq \sqrt[k]{\frac{1}{2}} \prod_{u=1}^t \lambda_u^{\sum_{j=-N_u+1}^{-M_u-1} \{w_{k+n}-w_k\}} \geq 2^{-N_t-1/k}.$$

Here the last inequality follows from $\lambda_u < \lambda_1 = 2$.

Let $L = \max\{|l_i| : 1 \leq i \leq k\}$. Then for every $t > \max(t_0, L)$, $i \in \{1, \dots, k\}$ and $N_t \leq n \leq \frac{m_t}{L}$,

$$T^{(l_i n)'}(w) \geq 2^{-N_t-1/k}$$

and so

$$S^{n'}(w_1, w_2, \dots, w_k) \geq 2^{-kN_t-1}.$$

Which together with (2.4) implies that every $\tilde{w} \in \mathbb{X}^k$ and $t > \max(t_0, k, L)$

$$\sum_{n=N_t}^{m_t/L} S^{n'}(\tilde{w}) \geq \left(\frac{m_t}{L} - N_t\right) 2^{-kN_t-1} \geq \frac{1}{2}.$$

Therefore for every $\tilde{w} \in \mathbb{X}^k$,

$$\sum_{n=1}^{\infty} S^{n'}(\tilde{w}) \geq \sum_t \sum_{n=N_t}^{m_t/L} S^{n'}(\tilde{w}) = \infty.$$

By Hopf's theorem for non-singular transformations S is conservative. □

The next example is a continuous time flow such that all the times are zero type and power weakly mixing. In this Markov Chain example the flow preserves an infinite measure.

2.2. The Markov Chain Example. A Borel map $X \times \mathbb{R} \ni (x, t) \mapsto \phi_t(x)$ such that

$$\phi_t \phi_s = \phi_{t+s}$$

is called a *non-singular flow* on (X, \mathcal{B}) . Given a measure μ on X , the semi-flow $\{\phi_t\}_{t \in [0, \infty)}$ is called exact if

$$\cap_{t \geq 0} \phi_t^{-1} \mathcal{B} = \{\emptyset, X\} \text{ mod } \mu.$$

A measure preserving flow $(X, \mathcal{B}, \mu, \{\phi_t\})$ is a *K-flow* if it admits an exhaustive and exact factor. Clearly a natural extension of an exact semiflow is a K-flow.

The flow is called *Power Weakly Mixing* if for every $t_1, t_2, \dots, t_n \in \mathbb{R}$,

$$\phi_{t_1} \times \phi_{t_2} \times \dots \times \phi_{t_n}$$

is an ergodic transformation of $\left((S^{\mathbb{R}})^k, \mathcal{B}^{\otimes k}, \mu^{\otimes k}\right)$.

A function $p : [0, \infty) \rightarrow [0, 1]$ is a *Markovian Renewal Function* if there exists a countable state space, which will be denoted by S , Markov Chain $\{X_t\}_{t \in [0, \infty)}$ and a state $a \in S$ such that

$$P_{a,a}(t) := P(X_t = a | X_0 = a) = p(t).$$

We say that p is *aperiodic* if

$$\gcd \{n \in \mathbb{N} : p(n) \neq 0\} = 1,$$

and *null recurrent* if

$$\sum_{n=1}^{\infty} p(n) = \infty \text{ and } p(n) \xrightarrow{n \rightarrow \infty} 0.$$

Given a renewal function p the sequence $\{p(n)\}_{n \in \mathbb{N}}$ defines a renewal sequence for the discrete time Markov Chain $\{X_n\}_{n=0}^{\infty}$. Thus if p is aperiodic and null recurrent then the Markov chain $\{X_n\}_{n=0}^{\infty}$ is aperiodic and null recurrent. Hence there exists a stationary (σ -finite) measure $\tilde{\mu} \in \mathcal{M}(S)$. It follows that the measure

$$\mu(\{a\}) = \int_0^1 \left(\sum_{s \in S} P_{s,a}(t) \tilde{\mu}(\{s\}) \right) dt$$

is a stationary measure for $(P_t)_{t \in \mathbb{R}} = \left(\{P_{a_1, a_2}(t)\}_{a_1, a_2 \in S} \right)_{t \in \mathbb{R}}$. Finally let $\nu = P^\mu$ be the measure on $S^{\mathbb{R}}$ with finite dimensional distributions

$$\nu[x_{t_0} = s_0, x_{t_1} = s_1, \dots, x_{t_n} = s_n] = \mu(\{s_0\}) P_{s_0, s_1}(t_1 - t_0) \dots P_{s_{n-1}, s_n}(t_n - t_{n-1}),$$

for every $t_0 < t_1 < \dots < t_n$ and $s_0, s_1, \dots, s_n \in S$. The flow $\{\phi_t\}_{t \in \mathbb{R}}$ on $S^{\mathbb{R}}$ defined by

$$\phi_t w(s) = w(s + t)$$

is ν - measure preserving. It is the natural extension of the semiflow $\{\phi_t\}_{t \in [0, \infty)}$.

Theorem 8. *Let $p : [0, \infty) \rightarrow [0, 1]$ be an aperiodic and null recurrent Markov Renewal Function then the flow $(S^{\mathbb{R}}, \mathcal{B}, \nu, \{\phi_t\}_{t \in \mathbb{R}})$ is conservative, exact and zero-type. If in addition for every $t_1, t_2, \dots, t_n \in \mathbb{R}_+$,*

$$\sum_{n=1}^{\infty} \prod_{j=1}^k p_{a,a}^{(n)}(t_j) = \infty$$

then the flow is Power Weakly Mixing.

Proof. Since $\{X_t\}_{t \geq 0}$ is a null recurrent Markov chain the flow is conservative and zero-type.

First we show that the tail σ -field of $\{X_t\}_{t \geq 0}$ is trivial, hence the semiflow $\{\phi_t\}_{t \in [0, \infty)}$ is exact. Let $h > 0$ and observe that $\Upsilon = (S^{h\mathbb{N}}, \mathcal{B}_{S^{h\mathbb{N}}}, \nu|_{h\mathbb{N}}, \phi_h)$ is a factor of $\mathfrak{X} = (S^{[0, \infty)}, \mathcal{B}, \nu, \phi_h)$. Since the discrete time chain $\{X_{nh}\}_{n \in \mathbb{N}}$ is aperiodic and recurrent, it follows by [BF] that its tail σ -algebra is trivial.

Denote by $\mathcal{F}_h = \mathcal{B} \cap S^{[0, h]}$. Then it follows from the Markov property that given Υ , for every $\Lambda_0, \Lambda_1, \dots, \Lambda_n \in \mathcal{F}_h$, the sets

$$[w \in \Lambda_1], [\phi_h w \in \Lambda_2], \dots, [\phi_{nh} w \in \Lambda_n]$$

are independent. By Kolmogorov's zero-one law \mathfrak{X} is an exact non-singular extension of Υ in the sense of [AD]. Therefore, since Υ is exact, it follows by Proposition 4 in [AD] that \mathfrak{X} is exact. See also [Ios, Theorem 6]. Therefore since the flow $\{\phi_t\}_{t \in \mathbb{R}}$ is the natural extension of the semiflow, it is a K-flow.

It follows that for every $t_1, t_2, \dots, t_k \in \mathbb{R}$ the transformation $R = \phi_{t_1} \times \phi_{t_2} \times \dots \times \phi_{t_k}$ is K and in order to prove ergodicity of R it is enough to show conservativity which is a consequence of the fact that,

$$\sum_{n=1}^{\infty} \prod_{j=1}^k p_{a,a}^{(n)}(|t_j|) = \infty.$$

□

Example: Let

$$p(t) = \frac{1}{\log(e+t)}.$$

Since $p(t)$ satisfies the conditions of [Kin, Theorem 6.6, p.144, see also p.41] there exists a continuous time Markov Chain $\{X_t : t \in \mathbb{R}\}$ on a countable state space S such that for

some $a \in S$,

$$p_{a,a}(t) := P(X_t = a | X_0 = a) = p(t).$$

Since

$$p(n) = \frac{1}{\log(e+n)} \neq 0, \quad \sum_{n=1}^{\infty} p(n) = \infty$$

and for every $t_1, t_2, \dots, t_n > 0$,

$$\sum_{n=1}^{\infty} \prod_{j=1}^k p_{a,a}^{(n)}(t_j) = \sum_{n=1}^{\infty} \prod_{j=1}^k p_{a,a}(n \cdot t_j) = \sum_{n=1}^{\infty} \prod_{j=1}^k \frac{1}{\log(e + t_j \cdot n)} = \infty,$$

it satisfies the conditions of Theorem 8 and hence the Markov flow defined by the Markov Chain is conservative, zero type and power weakly mixing.

Acknowledgments. The author would like to thank his advisor Prof. Jon Aaronson for his time, patience and advice.

REFERENCES

- [Aar] J. Aaronson, An introduction to infinite ergodic theory, Amer. Math. Soc., Providence, R.I., 1997. MR 1450400 (99d:28025).
- [AD] Aaronson, Jon; Denker, Manfred On exact group extensions. Ergodic theory and harmonic analysis (Mumbai, 1999). Sankhyā Ser. A 62 (2000), no. 3, 339–349.
- [AFS] T. Adams, N. Friedman, and C.E. Silva, Rank one power weak mixing for nonsingular transformations, Ergod. Th. & Dynam. Systems 21 (2001), 1321–1332.
- [BF] Blackwell, David; Freedman, David The tail σ -field of a Markov chain and a theorem of Orey. Ann. Math. Statist. 35 1964 1291–1295.
- [CKM] J. Coquet, T. Kamae and M. Mendès France, Sur la mesure spectrale de certaines suites arithmétiques, Bull. Soc. Math. France 105 (1977), 369–384.
- [Da1] A. I. Danilenko, ‘Funny rank one weak mixing for nonsingular Abelian actions’, Israel J. Math. 121 (2001) 29–54.
- [Da2] A. I. Danilenko, Infinite rank one actions and nonsingular Chacon transformations, Illinois J. Math. 48 (2004), no. 3, 769–786.
- [DP] Alexandre I. Danilenko; Kyewon K. Park, Rank-one flows of transformations with infinite ergodic index, Proc. Amer. Math. Soc. 139 (2011), 201–207.
- [DGSM] S. Day, B. Grivna, E. McCartney, and C.E. Silva, Power Weakly Mixing Infinite Transformations, New York J. of Math. 5 (1999), 17–24.
- [HK] A. Hajian and S. Kakutani, Weakly wandering sets and invariant measures, Trans. Amer. Math. Soc. 110 (1964), pp 136–151. MR 27:4904
- [Ham] T. Hamachi, On a Bernoulli shift with non-identical factor measures, Ergod. Th. & Dynam. Sys. 1 (1981), pp 273–283, MR 83i:28025, Zbl 597.28022.

- [Hel] E. Hellinger, Neue Begründung der Theorie quadratischer Formen von unendlich vielen Veränderlichen. *Journal für die reine and angewandte Mathematik* 36 (1909), pp 210–271.
- [Ios] Iosifescu, M. On finite tail σ -algebras. *Z. Wahrscheinlichkeitstheorie* 24 (1972), 159–166
- [Kin] Kingman, J. F. C. (1972). *Regenerative Phenomena*. John Wiley and Sons Ltd. ISBN 0-471-47905-5.
- [Kak] S. Kakutani, On equivalence of infinite product measures, *Ann. Math.* 49 (1948), pp 214–224.
- [Kos] Z. Kosloff, On a type III_1 Bernoulli shift, preprint arXiv:0912.1094.
- [Kre] U. Krengel, Transformations without finite invariant measures have finite strong generators, *Contributions to Ergodic Theory and Probability* (Ohio State Univ., Columbus, OH, 1970), *Lecture Notes in Math.*, no. 160, Springer-verlag, Berlin, 1970, pp. 135–157, MR 42 #4703, Zbl 201.38303.
- [KrS] U. Krengel and L. Sucheston, On mixing in infinite measure spaces. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete* 13 (1969), 150–164.
- [RS] Rosinski, J. and Samorodnitsky, G. (1996). Classes of mixing stable processes. *Bernoulli* 2 365–377. MR1440274
- [Ro] E. Roy. Ergodic properties of Poissonian ID processes. *Ann. Probab.*, 35(2):551–576, 2007.
- [ST] C. E. Silva and P. Thieullen, A skew product entropy for nonsingular transformations, *J. Lon. Math. Soc.* (2) 52 (1995), 497–516.